


Differentiable Maps Between Manifolds

Def. Let M and N be differentiable manifolds. A continuous function, $f: M \rightarrow N$, is said to be differentiable if for any coordinate chart $y: V \rightarrow \mathbb{R}^m$ on N and any chart $x: U \rightarrow \mathbb{R}^n$ on M , the map: $y \circ f \circ x^{-1}: x(U \cap f^{-1}(V)) \subseteq \mathbb{R}^n \rightarrow y(V) \subseteq \mathbb{R}^m$ is differentiable.



In practice how do we differentiate a function with respect to different coordinate charts?

Ex. Consider the unit sphere with the atlas defined by stereographic projections: $A = \{(S^2 - (0, 0, 1), \pi_N), (S^2 - (0, 0, -1), \pi_S)\}$ given

by: $(u, v) = \pi_N(x, y, z) = \left(\frac{x}{1-z}, \frac{y}{1-z} \right)$

$$(\bar{u}, \bar{v}) = \pi_S(x, y, z) = \left(\frac{x}{1+z}, \frac{-y}{1+z} \right)$$

Take the function $f: S^2 \rightarrow \mathbb{R}$ by $f(x, y, z) = z$.

- a. Find $\frac{\partial f}{\partial u}, \frac{\partial f}{\partial v}, \frac{\partial f}{\partial \bar{u}}, \frac{\partial f}{\partial \bar{v}}$
- b. Find formulas that relate $\frac{\partial f}{\partial u}$ to $\frac{\partial f}{\partial \bar{u}}$ and $\frac{\partial f}{\partial v}$, and $\frac{\partial f}{\partial v}$ to $\frac{\partial f}{\partial \bar{u}}$ and $\frac{\partial f}{\partial \bar{v}}$.
- c. Consider the point on the sphere in Cartesian coordinates given by $\left(\frac{1}{2}, \frac{\sqrt{2}}{2}, \frac{1}{2}\right)$. Find the coordinates in (u, v) and (\bar{u}, \bar{v}) .
- d. Show that the relationship in part b works for the point $\left(\frac{1}{2}, \frac{\sqrt{2}}{2}, \frac{1}{2}\right)$.

- a. Since $u = \frac{x}{1-z}$ and $v = \frac{y}{1-z}$ from $(u, v) = \pi_N(x, y, z) = \left(\frac{x}{1-z}, \frac{y}{1-z} \right)$

$$\pi_N^{-1}(u, v) = \left(\frac{2u}{u^2+v^2+1}, \frac{2v}{u^2+v^2+1}, \frac{u^2+v^2-1}{u^2+v^2+1} \right)$$

(from an earlier homework assignment)

That is

$$x = \frac{2u}{u^2+v^2+1}$$

$$y = \frac{2v}{u^2+v^2+1}$$

$$z = \frac{u^2+v^2-1}{u^2+v^2+1}.$$

By the chain rule:

$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial u}$$

(Or you can just substitute $z = \frac{u^2+v^2-1}{u^2+v^2+1}$ into f to get: $f(u, v) = \frac{u^2+v^2-1}{u^2+v^2+1}$).

We know $\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} = 0$ and $\frac{\partial f}{\partial z} = 1$, so we can write:

$$\frac{\partial f}{\partial u} = \frac{\partial z}{\partial u} = \frac{4u}{(1+u^2+v^2)^2}.$$

Similarly:

$$\begin{aligned}\frac{\partial f}{\partial v} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial v} = \frac{\partial z}{\partial v} \\ \frac{\partial f}{\partial v} &= \frac{4v}{(1+u^2+v^2)^2}.\end{aligned}$$

Let's take a similar approach to calculate $\frac{\partial f}{\partial \bar{u}}, \frac{\partial f}{\partial \bar{v}}$.

$$\begin{aligned}(\bar{u}, \bar{v}) &= \pi_S(x, y, z) = \left(\frac{x}{1+z}, \frac{-y}{1+z}\right) \\ \text{so } \bar{u} &= \frac{x}{1+z}, \quad \bar{v} = \frac{-y}{1+z}.\end{aligned}$$

We need to invert π_S .

Notice:

$$1 + (\bar{u})^2 + (\bar{v})^2 = \frac{x^2 + y^2 + (1+z)^2}{(1+z)^2} = \frac{2(1+z)}{(1+z)^2} = \frac{2}{1+z}$$

$$\text{Thus } 1 + z = \frac{2}{1 + (\bar{u})^2 + (\bar{v})^2} \quad \text{and } z = \frac{1 - (\bar{u})^2 - (\bar{v})^2}{1 + (\bar{u})^2 + (\bar{v})^2}.$$

$$x = \bar{u}(1 + z) = \frac{2\bar{u}}{1 + (\bar{u})^2 + (\bar{v})^2}$$

$$y = -\bar{v}(1 + z) = \frac{-2\bar{v}}{1 + (\bar{u})^2 + (\bar{v})^2}.$$

So we have:

$$\pi_S^{-1}(\bar{u}, \bar{v}) = \left(\frac{2\bar{u}}{1 + (\bar{u})^2 + (\bar{v})^2}, \frac{-2\bar{v}}{1 + (\bar{u})^2 + (\bar{v})^2}, \frac{1 - (\bar{u})^2 - (\bar{v})^2}{1 + (\bar{u})^2 + (\bar{v})^2} \right)$$

$$x = \frac{2\bar{u}}{1 + (\bar{u})^2 + (\bar{v})^2}; \quad y = \frac{-2\bar{v}}{1 + (\bar{u})^2 + (\bar{v})^2};$$

$$z = \frac{1 - (\bar{u})^2 - (\bar{v})^2}{1 + (\bar{u})^2 + (\bar{v})^2} = \frac{2}{1 + (\bar{u})^2 + (\bar{v})^2} - 1.$$

By the chain rule:

$$\frac{\partial f}{\partial \bar{u}} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial \bar{u}} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial \bar{u}} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial \bar{u}}$$

$$\frac{\partial f}{\partial \bar{v}} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial \bar{v}} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial \bar{v}} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial \bar{v}}.$$

Again, $\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} = 0$ and $\frac{\partial f}{\partial z} = 1$ so $\frac{\partial f}{\partial \bar{u}} = \frac{\partial z}{\partial \bar{u}}$, $\frac{\partial f}{\partial \bar{v}} = \frac{\partial z}{\partial \bar{v}}$.

Now we can say:

$$\frac{\partial f}{\partial \bar{u}} = \frac{-4\bar{u}}{(1+(\bar{u})^2+(\bar{v})^2)^2}, \quad \frac{\partial f}{\partial \bar{v}} = \frac{-4\bar{v}}{(1+(\bar{u})^2+(\bar{v})^2)^2}.$$

b. How do $\frac{\partial f}{\partial u}$, $\frac{\partial f}{\partial v}$, $\frac{\partial f}{\partial \bar{u}}$, and $\frac{\partial f}{\partial \bar{v}}$ relate to each other? Again it's by the chain rule.

$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial \bar{u}} \frac{\partial \bar{u}}{\partial u} + \frac{\partial f}{\partial \bar{v}} \frac{\partial \bar{v}}{\partial u}$$

$$\frac{\partial f}{\partial v} = \frac{\partial f}{\partial \bar{u}} \frac{\partial \bar{u}}{\partial v} + \frac{\partial f}{\partial \bar{v}} \frac{\partial \bar{v}}{\partial v}.$$

But how do we calculate $\frac{\partial \bar{u}}{\partial u}$, $\frac{\partial \bar{v}}{\partial u}$, etc?

u, v, \bar{u}, \bar{v} are related to each other by the transition function, that is:

$$\pi_S \circ \pi_N^{-1}(u, v) = (\bar{u}, \bar{v})$$

where: $\pi_N^{-1}(u, v) = \left(\frac{2u}{u^2+v^2+1}, \frac{2v}{u^2+v^2+1}, \frac{u^2+v^2-1}{u^2+v^2+1} \right)$

$$\pi_S(x, y, z) = \left(\frac{x}{1+z}, -\frac{y}{1+z} \right).$$

By a direct calculation we get:

$$(\bar{u}, \bar{v}) = \pi_S \circ \pi_N^{-1}(u, v) = \left(\frac{u}{u^2+v^2}, \frac{-v}{u^2+v^2} \right)$$

$$\bar{u} = \frac{u}{u^2+v^2} \quad \bar{v} = \frac{-v}{u^2+v^2}$$

$$\frac{\partial \bar{u}}{\partial u} = \frac{v^2-u^2}{(u^2+v^2)^2} \quad \frac{\partial \bar{u}}{\partial v} = \frac{-2uv}{(u^2+v^2)^2}$$

$$\frac{\partial \bar{v}}{\partial u} = \frac{2uv}{(u^2+v^2)^2} \quad \frac{\partial \bar{v}}{\partial v} = \frac{v^2-u^2}{(u^2+v^2)^2}$$

$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial \bar{u}} \frac{\partial \bar{u}}{\partial u} + \frac{\partial f}{\partial \bar{v}} \frac{\partial \bar{v}}{\partial u} = \frac{\partial f}{\partial \bar{u}} \left(\frac{v^2-u^2}{(u^2+v^2)^2} \right) + \frac{\partial f}{\partial \bar{v}} \left(\frac{2uv}{(u^2+v^2)^2} \right)$$

$$\frac{\partial f}{\partial v} = \frac{\partial f}{\partial \bar{u}} \frac{\partial \bar{u}}{\partial v} + \frac{\partial f}{\partial \bar{v}} \frac{\partial \bar{v}}{\partial v} = \frac{\partial f}{\partial \bar{u}} \left(\frac{-2uv}{(u^2+v^2)^2} \right) + \frac{\partial f}{\partial \bar{v}} \left(\frac{v^2-u^2}{(u^2+v^2)^2} \right).$$

c. For $(x, y, z) = \left(\frac{1}{2}, \frac{\sqrt{2}}{2}, \frac{1}{2} \right)$ in the coordinate system u, v we get:

$$u = \frac{x}{1-z} = \frac{\frac{1}{2}}{1-\frac{1}{2}} = 1 \quad v = \frac{y}{1-z} = \frac{\frac{\sqrt{2}}{2}}{1-\frac{1}{2}} = \sqrt{2}$$

$$\text{So } (u, v) = (1, \sqrt{2}).$$

In the coordinate system \bar{u}, \bar{v} we get:

$$\bar{u} = \frac{x}{1+z} = \frac{\frac{1}{2}}{1+\frac{1}{2}} = \frac{1}{3}; \quad \bar{v} = \frac{-y}{1+z} = \frac{-\frac{\sqrt{2}}{2}}{1+\frac{1}{2}} = -\frac{\sqrt{2}}{3}$$

$$\text{So } (\bar{u}, \bar{v}) = \left(\frac{1}{3}, -\frac{\sqrt{2}}{3} \right).$$

$$\begin{aligned}
 \text{d. } \frac{\partial f}{\partial u} &= \frac{4u}{(1+u^2+v^2)^2} & \text{at } (1, \sqrt{2}): \quad \frac{\partial f}{\partial u} &= \frac{4}{(1+1+2)^2} = \frac{1}{4} \\
 \frac{\partial f}{\partial v} &= \frac{4v}{(1+u^2+v^2)^2} & \text{at } (1, \sqrt{2}): \quad \frac{\partial f}{\partial v} &= \frac{4\sqrt{2}}{(4)^2} = \frac{\sqrt{2}}{4} \\
 \frac{\partial f}{\partial \bar{u}} &= \frac{-4\bar{u}}{(1+(\bar{u})^2+(\bar{v})^2)^2} & \text{at } \left(\frac{1}{3}, -\frac{\sqrt{2}}{3}\right): \quad \frac{\partial f}{\partial \bar{u}} &= \frac{-\frac{4}{3}}{\left(1+\frac{1}{9}+\frac{2}{9}\right)^2} = -\frac{3}{4} \\
 \frac{\partial f}{\partial \bar{v}} &= \frac{-4\bar{v}}{(1+(\bar{u})^2+(\bar{v})^2)^2} & \text{at } \left(\frac{1}{3}, -\frac{\sqrt{2}}{3}\right): \quad \frac{\partial f}{\partial \bar{v}} &= \frac{\frac{4\sqrt{2}}{3}}{\left(\frac{4}{3}\right)^2} = \frac{3\sqrt{2}}{4}.
 \end{aligned}$$

With $(u, v) = (1, \sqrt{2})$ and $(\bar{u}, \bar{v}) = \left(\frac{1}{3}, -\frac{\sqrt{2}}{3}\right)$ we can now directly check that:

$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial \bar{u}} \left(\frac{v^2 - u^2}{(u^2 + v^2)^2} \right) + \frac{\partial f}{\partial \bar{v}} \left(\frac{2uv}{(u^2 + v^2)^2} \right) = \left(-\frac{3}{4}\right) \left(\frac{2-1}{9}\right) + \left(\frac{3\sqrt{2}}{4}\right) \left(\frac{2\sqrt{2}}{9}\right) = \frac{1}{4}$$

$$\frac{\partial f}{\partial v} = \frac{\partial f}{\partial \bar{u}} \left(\frac{-2uv}{(u^2 + v^2)^2} \right) + \frac{\partial f}{\partial \bar{v}} \left(\frac{v^2 - u^2}{(u^2 + v^2)^2} \right) = \left(-\frac{3}{4}\right) \left(\frac{-2\sqrt{2}}{9}\right) + \left(\frac{3\sqrt{2}}{4}\right) \left(\frac{1}{9}\right) = \frac{\sqrt{2}}{4}.$$